



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07F 9/38, C07C 229/16, A61K 51/10, 51/08	A1	(11) International Publication Number: WO 00/24751 (43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/EP99/08031 (22) International Filing Date: 22 October 1999 (22.10.99) (30) Priority Data: 98402648.4 23 October 1998 (23.10.98) EP (71) Applicant (for all designated States except US): I.N.S.E.R.M. (INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE) [FR/FR]; 101, rue de Tolbiac, F-75654 Paris Cedex 13 (FR). (72) Inventors; and (75) Inventors/Applicants (for US only): GESTIN, Jean-François [FR/FR]; 5, chemin de la Coulée, F-44470 Mauves-sur-Loire (FR). LOUSSOUARN, Anthony [FR/FR]; 53, rue Fauré, F-44000 Nantes (FR). FAIVRE-CHAUVET, Alain [FR/FR]; 24, rue E. Zola, F-44300 Reze (FR). (74) Agents: DEMACHY, Charles et al.; Grosset-Fournier & Demachy, 20, rue de Maubeuge, F-75009 Paris (FR).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: CHELATING AGENTS FOR RADIOIMMUNOTHERAPY (57) Abstract <p>The invention relates to compounds of formula (I): in which: n_1, n_2, n_3 and n_4, represent an integer from 1 to 5, R_1, R_2, R_3 and R_4, independently from each other, represent $-\text{COOH}$, $-\text{PO}(\text{OH})_2$, at least one of R_1, R_2, R_3 or R_4 represents a group (II), wherein n_5 represents an integer from 1 to 5, R_5 represents $-\text{COOH}$ or $-\text{PO}(\text{OH})_2$, and Y represents H or a group $-(\text{CH}_2)_{n_6}-R_6$ in which n_6 represents an integer from 1 to 5, and R_6 represents $-\text{COOH}$ or $-\text{PO}(\text{OH})_2$, R represents H, or a group carrying a function linked to molecules able to bind with epitopes at the surface of cells. The invention also relates to the processes of preparation of said compounds, and to their use in pharmaceutical compositions and in diagnosis methods.</p> <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> <div style="text-align: center;"> <p style="text-align: right;">(II)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

CHELATING AGENTS FOR RADIOIMMUNOTHERAPY

5 The invention relates to compounds useful as chelating agents, complexes between said compounds and radioelements, and to their uses, in particular in pharmaceutical compositions and compositions for the diagnosis of pathologies such as cancers.

10 Immunotherapy with radiolabeled antibodies should allow fairly specific targeting of certain cancers (Schubiger et al., 1996; Parker, 1990). However, iodine-131 (Bardies et al., 1992; Stein et al., 1995) may not be the best isotope for tumor therapy because of its limited specific activity, low beta-energy, relatively long half-life and strong gamma emission.

15 Another approach to improving therapeutic efficacy is the use of replacement isotopes with better physical properties. Chelators that can hold radiometals with high stability under physiological conditions are essential to avoid excessive radiation damage to non-target cells. Moreover, the development of new bifunctional chelating agents is essential for this purpose. Thus synthesis of new chelating agents able to bind radiometals such as
20 rhodium-188, yttrium-90, samarium-153 or Bismuth-213 and in general all the α and β particles emitters will be required to possess sufficiently stable chelators.

 Accordingly, one of the aim of the invention is to provide chelating agents forming stable complexes *in vivo* with the numerous potential candidates for such applications.


25 The stability of a non-macrocyclic ligand can be favourably influenced by the preorganization of the open chain. In fact, a semi-rigid structure such as that of *trans* 1-2 diaminocyclohexane limits the rotation of the ethylene bridge, so that the purpose of the cyclohexane design is to preorient the four pendent arms in a skew position.

30 A first investigation (Mease et al., 1990), which was guided by a study performed on polyaminocarboxylic acid ligands incorporating the skeleton of ethylenediaminetetraacetic acid (EDTA) in a cyclohexane structure, showed the influence of this semi-rigid structure on the stability of the resulting complexes. A second study (Goeckeler et al., 1987) of the stability of lanthanides as ^{153}Sm -polyaminophosphonic acid complexes showed that ethylenediamine
35 tetramethylphosphonic acid (EDTMP) derivatives allow stable quantitative ^{153}Sm chelation.

The (1*R**, 2*R**, 4*S**)-4-acetamido-1,2-diaminocyclohexane dihydrochloride compound, the structural derivative of *trans*-1,2-diaminocyclohexane, have been prepared (Gestin et al., 1997; Loussouarn, et al., 1998). This intermediate, which is functionalized at position 4 of the cycle by a protected amine termination (Meares et al., 1984) for future covalent attachment to biomolecules, allows the introduction of different chelating groups via the free amines.

The Inventors have developed a new simple and efficient synthesis pathway from *trans*-1,2-diaminocyclohexane to provide access to a new class of semi-rigid chelating agents. This same reactional scheme applies to the reactional intermediary, (1*R**, 2*R**, 4*S**)-4-acetamido-1,2-diaminocyclohexane dihydrochloride, which allows the synthesis of these same chelating agents, though functionalized back of the cycle by a termination allowed coupling to an antibody or any other biological substance such as a hapten.

The present invention relates to compounds of the following formula (I) :



in which :

- n_1 , n_2 , n_3 and n_4 , independently from each other, represent an integer from 1 to 5, preferably from 1 to 3,

- R_1 , R_2 , R_3 and R_4 , independently from each other, represent :


. -COOH,

. -PO(OH)₂,

. -N $\begin{matrix} \diagup (CH_2)_{n_5}-R_5 \\ \diagdown Y \end{matrix}$

wherein n_5 represents an integer from 1 to 5, preferably from 1 to 3, R_5 represents -COOH or -PO(OH)₂, and Y represents H or a group -(CH₂) _{n_6} - R_6 in which n_6 represents an integer from 1 to 5, preferably from 1 to 3, and R_6 represents -COOH or -PO(OH)₂,

provided that at least one of R₁, R₂, R₃ or R₄ represents a group



such as defined above,



- R represents :

. H, or -NHCOCH₃, or

. a group carrying a function liable to bind, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, which are able to bind specifically with epitopes located at the surface of the cells of the organism, or to chemical or biological compounds located at the surface of a solid carrier, or

. a group carrying a function linked, if necessary via a binding site, to molecules as defined above,


the two following compounds, CDTPA and CTTHA, being excluded :



The invention relates more particularly to compounds of formula (I) such as defined above, characterized in that :

- when R₁, R₂, R₃ or R₄ represents -COOH or -PO(OH)₂, then n₁, n₂, n₃ or n₄ represents 1 respectively,


- when R₁, R₂, R₃ or R₄ represents a group



then n₁, n₂, n₃ or n₄ represents 2 or 3 respectively, and preferably 2,


- n₅, and optionally n₆, represents 1.

The invention also relates more particularly to compounds of formula (I) such as defined above, characterized in that at least one, and more preferably two of R₁, R₂, R₃ and R₄, represent a group



wherein n₅, n₆, R₅ and R₆ are defined above.

Preferred compounds of formula (I) such as defined above, wherein R is different from hydrogen, are compounds of the following formula (II) :



wherein n₁, n₂, n₃, n₄, R₁, R₂, R₃, R₄ and R are such as defined above.

The invention relates more particularly to compounds of formula (I) or (II) such as defined above, characterized in that R represents a group carrying a function liable to bind, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, as defined above, and in particular R represents a group chosen among all the coupling functions for vector or solid support binding, such as the following groups :


- . alcohol group, such as -OH,
- . amino group, such as -NH₂, -NO₂,
- . aldehyde group, such as -CHO,
- . carboxylic group, such as -COOH,
- . anhydride group, such as -CO-O-CO-R",
- . -CO-CH₂X, wherein X represents an halogen atom, such as Cl or Br,
- . -CO-X, wherein X represents an halogen atom, such as Cl or Br,
- . a diazonium ion N₂⁺,
- . an activated ester, such as -COOR", R" = ethyl or N-hydrosuccinimide,
- . a sulfonic group, such as SO₃H,
- . a thiocyanate group, such as -NCS, or an isocyanate -NCO, or a -NH-NCS group
- . a thiol group, such as -SH,
- . a disulfure group, such as -S-S-R".

The invention also concerns compounds of formula (I) or (II) such as defined above, characterized in that R represents a group carrying a function linked, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, as defined above, and more particularly R represents a group chosen among the following groups :

- . -O-CO-R',
- . -NH-CO-R',
- . -NH-CS-R',
- . -CH=N-R',
- . -CO-NH-R',
- . -CO-CH₂-NH-R',
- . -N=N-NH-R',
- . -SO₂-NH-R',
- . -NH-CS-NH-R',
- . -thioether-R',
- . -CO-S-R',
- . -CO-CH₂-S-R',
- . -S-S-R',
- . -NH-CH₂-R',
- . -CO-NH-N=CH-R',
- . -CS-NH-N=CH-R',

wherein R' represents said molecule.

The invention concerns more specifically compounds such as described above of the following formula (III) :




in which R₁, R₂, R₃, R₅ and R₆ independently from each other represent -COOH or -PO(OH)₂, and R is a group as defined above.

Preferred compounds of formula (III) are such that :

- . R₁ = R₅ = R₆ = COOH and R₂, = R₃ = PO(HO)₂, or
- . R₁ = R₂ = R₃ = R₅ = R₆ = COOH, or
- . R₁ = R₂ = R₃ = R₅ = R₆ = PO(OH)₂.

The invention also concerns more specifically compounds such as described above, of the following formula (IV) :



wherein R_2 , R_5 and R_6 , independently from each other, represent $-\text{COOH}$ or $-\text{PO}(\text{OH})_2$, and R is a group as defined above.

Preferred compounds of formula (IV) are such that :

- . $R_2 = R_3 = \text{PO}(\text{OH})_2$, and $R_5 = R_6 = \text{COOH}$, or
- . $R_2 = R_3 = R_5 = R_6 = \text{COOH}$, or
- . $R_2 = R_3 = R_5 = R_6 = \text{PO}(\text{OH})_2$.

The invention also relates to complexes between a compound such as described above, and a radioactive element, said complexes resulting from the association of said radioelement with the $-\text{COOH}$ and/or $-\text{PO}(\text{OH})_2$ groups of said compound (the bonds between said radioelement and said compound being ionic bonds).

The above-mentioned radioelements are more particularly α , β or γ emitter radiometals, and preferably from the groups of actinides or lanthanides.

The invention relates more particularly to complexes such as described above, characterized in that said radioelements are α or β emitter radiometals (susceptible to be used in therapy, and more particularly in radioimmunotherapy in the frame of cancer treatments).

Advantageously, α emitter radiometals are chosen among the followings : Actinium 225, Bismuth 213.

Advantageously, β emitter radiometals are chosen among the followings : ^{33}P , ^{199}Au , ^{121}Sn , ^{177}Lu , ^{67}Cu , ^{105}Rh , ^{47}Sc , ^{77}As , ^{153}Sm , ^{159}Gd , ^{143}Pr , ^{186}Re , ^{111}Ag , ^{149}Pm , ^{109}Pd , ^{166}Ho , ^{32}P , ^{188}Re , ^{194}Ir , ^{142}Pr , ^{90}Y .

Preferred complexes with radiometals used in therapy, as defined above, are such that the compound is chosen among those wherein R represents a group carrying a function linked, if necessary via a binding site, to molecules as defined above, and more particularly among those compounds wherein the group R comprises :

- an antibody (polyclonal or monoclonal) liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism,

- or an hapten, i.e. a non-immunogenic molecule of low MW capable of inducing the production of antibodies against itself, said hapten being liable to recognize and to bind to one or several molecules already bound (in a first step of the treatment) to epitopes on the surface of specific cells in the organism,

- or a peptide resulting from the association of different amino acids and liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism.

The invention also concerns the use of a complex such as described above, for the manufacture of a medicament for radioimmunotherapy (also called radiopharmaceutical), in particular for the treatment of cancers, or for the treatment against metastase proliferation.

More particularly, the invention relates to the use of a complex such as defined above, for the manufacture of a medicament for the treatment of :

- lung cancer, said complex preferably being such that it comprises a radioelement chosen among : ^{188}Re , ^{186}Re , ^{153}Sm , ^{67}Cu and ^{90}Y , and wherein R comprises an antibody specific for lung cancer cells, such as Anti N-CAM Antibody, Anti CEA Antibody, Anti Carbohydrates Antibodies, or an hapten chosen among Anti N-CAM-679 Bispecific antibody, Anti CEA-679 Bispecific antibody, Anti Carbohydrates-679 Bispecific antibody, Anti N-CAM-734 Bispecific antibody, Anti CEA-734 Bispecific antibody, Anti Carbohydrates-734 Bispecific antibody,

- liver and pancreatic cancers, said complex preferably being such that it comprises a radioelement chosen among those cited above in the case of lung cancer, and wherein R comprises an antibody specific for liver and pancreatic cancer cells, such as antibodies and haptens described above in the case of lung cancer,

- ovarian cancer, said complex preferably being such that it comprises a radioelement chosen among those cited above in the case of lung cancer, and wherein R comprises an antibody specific for ovarian cancer cells, such as OC125, MOV18, MOV19, OVTL3, or an hapten chosen among OC125-679 Bispecific antibody, MOV18-679 Bispecific antibody, MOV19-679 Bispecific antibody, OVTL3-679 Bispecific antibody, OC125-734 Bispecific antibody, MOV18-734 Bispecific antibody, MOV19-734 Bispecific antibody, OVTL3-734 Bispecific antibody,

- bladder cancer, said complex preferably being such that it comprises a radioelement chosen among those cited above in the case of lung cancer, and

wherein R comprises an antibody specific for bladder cancer cells, such as AC48-127, or an hapten chosen among 48-127 Bispecific antibody, 48-127-679 Bispecific antibody, 48-127-734 Bispecific antibody,

- colorectal cancer, said complex preferably being such that it comprises a radioelement chosen among those cited above in the case of lung cancer, and wherein R comprises an antibody specific for colorectal cancer cells, such as Anti CEA Antibody, Anti Carbohydrates Antibodies, or an hapten chosen among Anti CEA-679 Bispecific antibody, Anti Carbohydrates-679 Bispecific antibody, Anti CEA-734 Bispecific antibody, Anti Carbohydrates-734 Bispecific antibody,

- thyroid medullary cancer, said complex preferably being such that it comprises a radioelement chosen among those cited above in the case of lung cancer, and wherein R comprises an antibody specific for thyroid medullary cancer cells, such as Anti CEA Antibody, or an hapten chosen among Anti CEA-679 Bispecific antibody, Anti CEA-734 Bispecific antibody,

- lymphoma, said complex preferably being such that it comprises a radioelement chosen among : ^{213}Bi , ^{225}Ac , ^{153}Sm , and ^{67}Cu , and wherein R comprises an antibody specific for lymphoma cells, such as specific antibody against expressed antigens surfaces lymphocyte cells, e.g. CD19, CD37, or an hapten such as bispecific antibody against expressed antigens surfaces lymphocyte cells, e.g. CD19-679, CD37-679, CD19-734, CD37-734,

- myeloma, said complex preferably being such that it comprises a radioelement chosen among : ^{213}Bi , ^{225}Ac , ^{153}Sm , and ^{67}Cu , and wherein R comprises an antibody specific for myeloma cells, such as specific antibody against expressed antigens surfaces myeloma cells, e.g. BB4, or an hapten such as bispecific antibody against expressed antigens surfaces myeloma cells, BB4-679, BB4-734,

- osteoarticular pathology, particularly in bone cancer extension balance.

The invention also concerns pharmaceutical compositions characterized in that they comprise an effective amount of a complex such as described above, in association with a suitable pharmaceutical carrier.

Pharmaceutical compositions according to the invention are more particularly characterized in that they are in a form suitable for an IV or IP administration in located areas.

Preferred pharmaceutical compositions according to the invention, are characterized in that the daily dosage is comprised between 1 and 100MBq /kg, e.g. between 3,7 and 74MBq/kg.

The invention also relates to complexes, as defined above, between a compound such as described above, and a radioactive element, characterized in that the radioelements are γ emitter radiometals (i.e. radiometals susceptible to be used in diagnosis methods, such as radioimmunosciintigraphy).

Advantageously, said radiometals are chosen among ^{111}In , $^{99\text{m}}\text{Tc}$, ^{64}Cu .

Preferred complexes with radiometals used in diagnosis, as defined above, are such that the compound is chosen among those wherein R represents a group carrying a function linked, if necessary via a binding site, to molecules as defined above, and more particularly among those compounds wherein the group R comprises :

- an antibody (polyclonal or monoclonal) liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism,

- or an hapten, i.e. a non-immunogenic molecule of low MW capable of inducing the production of antibodies against itself, said hapten being liable to recognize and to bind to one or several molecules already bound (in a first step of the method of diagnosis) to epitopes on the surface of specific cells in the organism,

- or a peptide resulting from the association of different amino acids and liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism.

The invention also concerns the use of a complex such as described above, for carrying out diagnosis methods such as radioimmunosciintigraphy.

More particularly, the invention concerns the use of a complex such as defined above, for carrying out the following diagnosis methods by radioimmunosciintigraphy :

- diagnosis of cancers, such as cited above, the complex used being preferably such that it comprises ^{111}In , or $^{99\text{m}}\text{Tc}$ as radioelements, and R comprises an antibody or a hapten specific for such cancer cells, such as antibodies or haptens mentioned above in the frame of the cancers listed above,

- diagnosis of cardiovascular diseases, such as graft rejection, myocardic infarcts,

- diagnosis of cerebral diseases,

- diagnosis of renal diseases, in particular in the study of individual kidney functions, location of ectopic kidney, renal filtration and secretion troubles,

- vascular diseases, such as embolism and thrombosis, the complex used being preferably such that it comprises ^{111}In , or $^{99\text{m}}\text{Tc}$ as radioelements, and R comprises an antibody such as anti platelets or anti fibrin antibodies.

The invention also concerns the use of the complexes defined above for carrying out bone scintigraphy, in particular in the frame of the diagnosis of osteoarticular pathology, particularly in bone cancer extension balance.

5 The invention also relates to methods for the *in vitro* diagnosis of pathologies listed above, characterized in that it comprises the steps of incubating a biological sample, such as serum, plasma urines, with a complex as described above, the components of the biological sample being fixed to a solid carrier, rinsing the solid carrier and detecting the γ emission of the complex bound to the components of the sample on the solid carrier.

10 The invention also concerns the kits for carrying out said diagnosis methods, said kits comprising complexes as described above according to the invention.

The invention also relates to the use of a compound of formula (I) defined above, included compounds CDTPA and CTTHA, for the manufacture of a medicament for the treatment of pathologies where ionic imbalances occur, or against the formation of stones in the organism.

15 The invention also relates to the use of a compound of formula (I) defined above, included compounds CDTPA and CTTHA, for carrying out a process for the detoxication of polluted medium, such as liquid phases polluted by bivalent or trivalent metals radioactives or not.

20 The invention also concerns a process for the detoxication of a polluted medium comprising the steps of contacting said medium with a compound as defined above, advantageously itself bound to a solid carrier, and recovering said medium substantially free of contaminants which are bound to said compound on the solid carrier.

25 The invention also relates to the use of a compound of formula (I) defined above, included compounds CDTPA and CTTHA, for carrying out a process for the radionuclides purification, said compound of the invention being bound to a solid phase.


30 The invention also relates to the use of a complex between of formula (I) defined above, included compounds CDTPA and CTTHA, for carrying out a bone scintigraphy, in particular in the frame of the diagnosis of osteoarticular pathology, particularly in bone cancer extension balance.

35 The invention also relates to processes for preparing compounds and complexes as described above.

A process for the preparation of compounds according to the invention, comprises the following steps :


The invention also relates to processes for preparing compounds and complexes as described above. A process for the preparation of compounds according to the invention, comprises the following steps :

- contacting trans-1,2-diaminocyclohexane of the following formula a :




wherein R_a is H or NHCOCH_3 ,


10 * either with vinyl propionate, preferably by stirring 20h at room temperature, leading to the following compound b



20 . contacting compound b with $\text{X}-(\text{CH}_2)_n-\text{COOEt}$, wherein X represents an halogen atom, and n represents an integer from 1 to 5, preferably at reflux during 15h, leading to the following compound c




25 . treating compound c with HCl, preferably 6N HCl at reflux overnight, leading to the following compound d




wherein R_b represents H or NH_2 ,


35 * or with diethyl vinyl phosphonate, preferably by stirring 15h at reflux, leading to the following compound e



. contacting compound e with $X-(CH_2)_n-COOEt$, wherein X represents an halogen atom, and n represents an integer from 1 to 5, preferably at reflux during 15h, leading to the following compound f




. treating compound f with HCl, preferably 6N HCl at reflux overnight, leading to the following compound g




wherein R_b represents H or NH_2 ,

. if desired, treating compound g with phosphorous acid, preferably by stirring 30 mn at $80^\circ C$, leading to the following compound h




Another process for the preparation of compounds according to the invention, comprises the following steps :


- contacting the compound of formula a described above with a compound of formula $H_2C=CH-(CH_2)_{nA}-CN$ wherein $nA = 0$ (acrylonitrile), or nA represents a integer from 1 to 3, preferably at room temperature during 20h, leading to the following compound i




- contacting compound i with the following compound j



wherein X represents an halogen atom, na and nb, independently from each other represent an integer from 1 to 5, Z represents H or (CH₂)_{nc}-COOEt, and nc represents an integer from 1 to 5, preferably at 70°C during 2 days, leading to the following compound k




. treating compound k with HCl, preferably 6N HCl at reflux overnight, leading to the following compound l




wherein Z_a represents H or $-(CH_2)_{n_b}-COOH$, n_a , n_a and n_b being such as defined above, and R_b represents H or NH_2 .

Another process for the preparation of compounds according to the invention, comprises the following steps :

- contacting the compound of formula a described above with paraformaldehyde and diethylphosphite, preferably in THF at reflux during 4h, leading to the following compound m




- treating compound m with HCl, preferably 3N HCl in MeOH at 50°C overnight, leading to the following compound n




- contacting compound n :


* either with 1 equivalent of compound j




15 as described above, preferably at 70°C during 2 days, leading to the following compound of formula o



25 . contacting compound o with X-(CH2)n-COOEt, wherein X represents an halogen atom, and n represents an integer from 1 to 5, preferably at reflux during 15h, leading to the following compound p




35 . treating compound p with HCl, preferably 6N HCl at reflux overnight, leading to the following compound q




wherein Z_a represents H or $-(CH_2)_{nb}-COOH$, n_a , n_b and n being such as defined above, R_b represents H or NH_2 ,

if desired, treating compound q with phosphorous acid, preferably by stirring 30 mn at $80^\circ C$, leading to the following compound r




wherein Z_b represents H or $-(CH_2)_{nb}-PO(OH)_2$


* or with 2 equivalents of compound j



as described above, preferably at $70^\circ C$ during 2 days, leading to the following compound of formula s




. treating compound s with HCl, preferably 6N HCl at reflux overnight, leading to the following compound t



wherein Za represents H or $-(CH_2)_{nb}-COOH$, na, nb and n being such as defined above, Rb represents H or NH_2 ,

. if desired, treating compound t with phosphorous acid, preferably by stirring 30 mn at $80^\circ C$, leading to the following compound u



wherein Zb represents H or $-(CH_2)_{nb}-PO(OH)_2$.

Compound of formula a can be obtained according to the method described in Gestin et al., 1997, and Loussouarn et al., 1998.

30 Compounds wherein Rb represents NH_2 obtained according to the processes described above, can then be transformed in order to correspond to compounds of formula (I) wherein R represents a group carrying a function liable to bind, if necessary via a binding site, to molecules as defined above.

35 By way of example, compounds of formula (I) wherein R represents $-N=C=S$, can be obtained by treatment of said compounds wherein Rb represents NH_2 with $CSCl_2$, preferably in acidic or basic conditions.

Compounds of formula (I) wherein R represents a group carrying a function linked, if necessary via a binding site, to molecules as defined above,

can then be obtained by coupling said compounds, wherein R represents a group carrying a function liable to bind to said molecules, with said molecules.


Complexes according to the invention are advantageously obtained by incubating the compounds with the radioelements at 37°C during 3 hours.

The invention will be further illustrated in the following examples for the preparation of compounds and complexes according to the invention.


In order to save the functionalized reactional intermediate, i.e. the (1*R**, 2*R**, 4*S**)-4-acetamido-1,2-diaminocyclohexane dihydro chloride compound, the commercial product, *trans*-1,2-diaminocyclohexane 1, has been used as starting material.

Two approach routes to differently substituted amines were carried out successfully.

- The first depicted in schemes I and II was the Michael type addition of primary amines to some vinylic derivatives to provide monoaddition with high selectivity (Bergeron et al., 1981), allowing N-alkylation to be envisaged at this step. In strategy depicted in scheme I, compounds 2a and 2b were alkylated by ethyl bromoacetate under conditions recommended by Studer (Studer and Meares, 1992) (KI and Na₂CO₃) to give tetraesters 3a and 3b. Acid-catalysed hydrolysis of the ester functions was performed in 3M hydrochloric acid to give the tetracarboxylic acid 4a and the mixed acid 4b. At last, in order to generate the structure 4c, carboxylic functions were converted into phosphonic functions using H₃PO₃/PCl₃ according to the method of Krüger and Bauer (Krüger and Bauer, 1972). The other strategy described in scheme II required preparation of protected bis-carboxymethylated amino ethyl bromide. In view to convenience of deprotecting ethyl esters by acid-catalysed hydrolysis, *N,N*-bis(ethylacetate)-2-bromoethyl-amine was prepared according to the Williams and Rapoport's procedure (Williams and Rapoport, 1994) with minor modifications. N-alkylation of 2d with the branching group in a mixed solvent system (CH₃CN/EtOH) at 70°C gave 3d in 60% yield. Acid-catalysed hydrolysis of the ester functions as well as nitriles is the more convenient method (Ornstein et al., 1989), of hexacarboxylic acid 4d preparation.




scheme I




scheme II

- In the case of certain diamines as *trans*-1,2-diaminocyclohexane **1**, the second route, as shown in schemes III, IV allowed the aminophosphonomethylation of amines protected by a methylene bridge between the two nitrogen atoms of **1**. This protecting group will subsequently provide for a different functionalization on the amine. The reaction of Kabachnick-field described and detailed by Baily and Burgada (Baily and Burgada, 1995), gave compound **6** which was prepared from paraformaldehyde and diethylphosphite in THF. **7** was obtained by removing the protecting group in acidic conditions.



scheme III


The monoalkylation or the dialkylation (see scheme IV), depending on the stoichiometry of the reaction gave respectively compounds **8** and **12**. The mixed acid **13** was obtained after hydrolysis of **12** in 6M hydrochloric acid and the hexaphosphonic acid **14** was prepared according to the method of Bauer and Kruger as described above. The synthesis of chelating agents **10** and **11** required an additional step which was the alkylation of **8** by ethyl bromoacetate to give the mixed ester **9**. Acid-catalysed hydrolysis gave **10** and **11** after reaction of conversion described all above.



Scheme IV

In conclusion, different non-functionalized ligands bearing aminophosphonate or aminocarboxylate chelate groups and mixed chelate groups were prepared and tested for their complexation properties with ¹⁵³Sm. The synthetic method described above was applied to the previously synthesized intermediate, the (1*R**, 2*R**, 4*S**)-4-acetamido-1,2-diaminocyclohexane, resulting in the synthesis of several polyaminocarboxylic acids, polyaminophosphonic acids and mixed semi-rigid functionalized ligands (BCA).

The different access routes to non-functionalized compounds described here were used without modifying the synthesis in order to obtain their functionalized homologues ready to be used in a coupling reaction as described in scheme V. We observed the influence of aminocarboxylic acid and aminophosphonic acid functions on the stability of the resulting complexes.



Scheme V

Experimental

General Procedures

All experiments were performed under nitrogen. Solvents were distilled prior to reactions. The primary chemicals used were commercial products (Sigma-Aldrich Company). Product purity and reaction progress were monitored on thin-layer chromatography (TLC) plates (60 F254, Merck), and liquid chromatography was carried out on a silica gel column (Merck 60,70-230 mesh). TLC revelation was performed under UV light (254 nm) or by iodine.

Nuclear Magnetic Resonance (MNR)

¹H and ¹³C NMR spectra were recorded on a Bruker AC 250 spectrometer (250 Mhz). Chemical shifts are reported in ppm to phosphoric acid as reference (85 % H₃PO₄ in heavy water), positive values being downfield.

Chemical shifts (δ) are reported in ppm. Coupling constant J is reported in Hertz (Hz).

Mass Spectrometry (MS)

MS spectra were recorded on a Mat Finnigan LCQ Ion Trap mass apparatus using the electrospray method in negative or positive mode.

Starting Material

The bisphosphonate **6** was prepared in our laboratory according the synthesis procedure of Baily and Burgada with minor modifications.

Synthesis and Spectroscopic data

N,N'-[(2-ethoxycarbonyl)eth-1-yl]-trans-cyclohexane-1,2-diamine 2a:

To freshly distilled *trans*-1,2-diaminocyclohexane **1** (1 ml, 8.33 mmol) in 50 ml of ethanol was added vinyl propionate (1.50 ml, 13.7 mmol) in one portion. After stirring 20h at room temperature, the reaction mixture was concentrated by rotary evaporation to yield a pale yellow oil (2.6 g, 8.32 mmol, 100%) which was used directly in the next step. ¹H NMR (CDCl₃): δ 1.22 (t, 12H), 1.67 (m, 2H), 1.82 (m, 2H), 2.06 (m, 2H+2H), 2.43 (t, 4H), 2.67 (dt, 2H), 2.98 (dt, 2H), 4.10 (q, 4H). ¹³C NMR (CDCl₃): δ 14.17, 24.31, 31.46, 35.34, 42.19, 60.23, 61.29, 172.69. (M+H⁺): 315

N,N'-[(2-diethylphosphono)eth-1-yl]-trans-cyclohexane-1,2-diamine 2b:

To freshly distilled *trans*-1,2-diaminocyclohexane **1** (1 ml, 8.33 mmol) in 50 ml of ethanol was added diethyl vinyl phosphonate (2.80 ml, 18.21 mmol). The reaction mixture was allowed to stir at reflux during 15 hours. After removal the solvent under reduced pressure, the resulting oil was purified by column chromatography (silica gel, CH₂Cl₂-EtOH 1:1) to give 2.9 g of a limpid oil (6.65 mmol, 80%). ¹H NMR (CDCl₃): δ 0.94 (m, 2H), 1.15 (m, 2H), 1.24 (t, 12H), 1.64 (m, 2H), 1.83-1.95 (m, 8H), 2.05 (m, 2H), 2.71 (dt, 2H), 2.97 (dt,

2H), 4.07 (dq, 8H). ^{13}C NMR (CDCl_3): d 16.37, 16.46, 25.42 (JC-P : 149 Hz), 28.20, 31.41, 40.47, 40.51, 61.29, 61.40, 61.50. (M+H⁺): 443

N,N'-[(2-cyano)eth-1-yl]-*trans*-cyclohexane-1,2-diamine 2d: To freshly
5 distilled *trans*-1,2-diaminocyclohexane 1 (1 ml, 8.33 mmol) in 50 ml of ethanol
was added acrylonitrile (1.20 ml, 18.32 mmol). After stirring 20h at room
temperature, the reaction mixture was concentrated by rotary evaporation to
yield an pale yellow oil witch was purified by recrystallisation in diethylether to
give 1.40 g of a white solid (6.35 mmol, 78%): mp : 65°C

10 ^1H NMR (CDCl_3): d 1.02 (m, 2H), 1.22 (m, 2H), 1.70-1.79 (m, 2H +
2H), 2.02-2.17 (m, 2H+2H), 2.49 (t, 4H), 2.80 (dt, 2H), 3.02 (dt, 2H). ^{13}C
NMR (CDCl_3): d (M+H⁺): 221

N,N'-[(2-ethoxycarbonyl)eth-1-yl]-*N,N'*-(ethylacetate)-*trans*-
15 cyclohexane-1,2-diamine 3a: To a solution of 2a (1 g; 3.18 mmol) in 50 ml of
freshly distilled CH_3CN under nitrogen were added Na_2CO_3 (0.50 g; 3.01
mmol) and KI (g; mmol). After stirring for 1 hour at 60°C, $\text{BrCH}_2\text{COOEt}$
(1.80 ml; 7.15 mmol) was added dropwise. The reaction mixture was kept at
this temperature over a period of 24 hours prior to cooling to room temperature,
20 filtration and concentration under reduced pressure. The residue was taken up in
 CHCl_3 (200 ml) and washed with water. The organic layer was dried (Na_2SO_4)
and concentrated under reduced pressure to give a yellow-brown oil. The crude
product was purified by column chromatography (silica gel, CH_2Cl_2 -EtOH
95:5). The fractions containing pure product were collected and dried to give an
25 limpid oil (0.8 g; 1.56 mmol; 49 %). ^1H NMR (CDCl_3): d 1.10 (m, 4H), 1.26
(t, 12H), 1.91 (m, 2H), 2.00 (m, 2H), 2.51 (dt+m, 4H+2H(CH cycle)), 2.95
(dt, 4H), 3.39 (d, 4H), 4.12 (m, 8H). (M+H⁺): 515

N,N'-(ethylacetate)-*N,N'*-[(2-diethylphosphono)eth-1-yl]-*trans*-
30 cyclohexane-1,2-diamine 3b: The tetraester has been prepared as described
above for compound from 1 g of compound 2b, Na_2CO_3 (0.70g, 6.60 mmol),
KI (0.40, 2.40 mmol) and ethylbromoacetate (1.80 ml; 7.15 mmol). Purification
by chromatography (SiO_2 (CH_2Cl_2 -MeOH 99 : 1) gave 0.58 g of a pale yellow
oil (0.94 mmol; 41%). ^1H NMR (CDCl_3): d 1.11 (m, 4H), 1.25 (t, 6H), 1.29 (t,
35 12H), 1.70 (m, 2H), 1.85-2.10 (m, 4H+2H), 2.90 (t, 4H), 3.38 (d, 4H), 4.09
(m, 12H). ^{13}C NMR (CDCl_3): d 14.11, 16.31, 16.41, 25.63, 26.03 (JC-P :
135), 27.98, 44.86, 52.43, 60.26, 61.31, 61.41, 63.08, 172.46. (M+H⁺): 615

General procedure for preparation of corresponding acids: *trans*-cyclohexane-1,2-diamine-*N,N'*-acetic-*N,N'*-propionic acid 4a and *trans*-cyclohexane-1,2-diamine-*N,N'*-acetic-*N,N'*-ethylphosphonic acid 4b:

Compound 3a or 3b (1 g) was dissolved in 6N aqueous hydrochloric acid (12ml) and heated to reflux overnight. The refrigerant was removed, and the reaction mixture was kept at 70°C to dryness. An additional aqueous hydrochloric acid 6N (12 ml) was then added, and the solution was heated to dryness. the residue was taken up in MeOH and evaporated under reduced pressure. This step repeated twice gave the corresponding acid as an off-white solid, which was dried under vacuum and kept under nitrogen.

Compound 4a: ¹H NMR (D₂O): d 1.25-1.40 (m, 4H), 1.60-2.15 (m, 4H), 2.28 (m, 2H), 2.75 (t, 2H), 2.96 (t, 2H), 3.22 (m, 2H), 3.50-3.90 (m, 4H), 4.15 (s, 1H), 4.28 (s, 1H) ¹³C NMR (CDCl₃): d 25.66, 26.22, 28.87, 31.20, 31.99, 47.25, 54.93, 66.92, 175.03, 176.46 (M-H⁺): 373

Compound 4b: ¹H NMR (D₂O): d 1.05-1.40 (m, 4H), 1.65-2.15 (m, 4H), 2.90-3.25 (m, 6H), 3.45-3.65(m, 2H), 3.70-3.95 (m, 2H). ¹³C NMR (CDCl₃): d 25.66, 26.22, 28.87, 31.20, 31.99, 47.25, 54.93, 66.33, 176.73 (M-H⁺): 445

***trans*-cyclohexane-1,2-diamine-*N,N'*-ethylphosphonic-*N,N'*-methylphosphonic acid 4c:** A mixture of compound 4c (0.5 g; 1.12 mmol) and phosphorous acid (0.202 g; 2.46 mmol) in 10 ml of dry toluene was heated to 80°C and stirred for 30 min. PCl₃ (0.22 ml; 2.46 mmol) was then added dropwise, and the reaction mixture was kept at this temperature for 20 hours before being cooled to room temperature. The solvent was discarded and the residual product dissolved in a small volume of water. After filtration, the filtrate was evaporated to give a residue which was purified by precipitation in warm acetone and collected by filtration. The purification step was repeated twice to give 4c which was dried under vacuum and kept under nitrogen (0.460 g; 0.83 mmol; 74%). ¹H NMR (D₂O): d 1.15-1.65 (m, 4H), 1.75-2.10 (m, 2H), 2.15-2.40 (m, 6H), 3.00-3.60 (m, 10H). ¹³C NMR (CDCl₃): d (M-H⁺): 517.

***N,N*-Bis(ethylacetate)-2-bromoethyl-amine 5:** Bromide 5 was synthesised in our laboratory according the synthesis procedure of Williams and Rapoport with minor modifications concerning the bis N-alkylated ethanolamine synthesis. To a 4°C solution of ethanolamine (6ml; 0.1 mol) in 100 ml of dried acetonitrile was added dropwise ethylbromoacetate (7.4 ml; 66 mmol) over a period of 20

min during which time a large quantity of precipitate formed. The mixture was allowed to stir for 2 hours at this temperature. The white solid was removed by filtration and washed with a small quantity of acetonitrile. The filtrate was concentrated under reduced pressure. The resulting liquid was taken up in CHCl₃ (100mL) and washed with water. The organic layer was dried (Na₂SO₄) and concentrated under reduced pressure to give a liquid which was used directly in the next step (5.9 g; 25.32 mmol; 77%). The dialkylated ethanolamine and Ph₃P (7.72 g; 27.9 mmol) were dissolved in CH₂Cl₂ (100mL). The mixture was cooled in an ice bath and vigorously stirred while NBS (4.96 g; 27.9 mmol) was added in small portions. After the solution was stirred at 0°C for two hours, evaporation of the solvent gave a semisolid which was triturated with ether and the resulting solid was separated by filtration. the filtrate was evaporated to give an oil which was purified by column chromatography (silica gel, CH₃Cl). (6.14 g; 20.7 mmol; 62% overall) ¹H NMR (CDCl₃): d 1.26 (m, 6H), 3.15 (t, 2H, J = 7.75 Hz), 3.44 (t, 2H, J = 7.75 Hz), 3.59 (s, 4H), 4.15 (q, 4H). ¹³C NMR (CDCl₃): d (M+H⁺): 297

N,N'-(2-cyano)eth-1-yl]-*N,N'*-[*N''*,*N''*'-bis-(ethylacetate-2-aminoethyl)]-*trans*-cyclohexane-1,2-diamine 3d: To a solution of compound 2d (1 g; 4.54 mmol) and bromide 5 (3 g; 10.13 mmol) in a mixed solvent system (CH₃CN-EtOH, 1:1) was added Na₂CO₃ (1.4 g; 13.20 mmol) and KI (0.75 g; 4.54 mmol). After stirring for 2 days at 70°C, the reaction mixture was filtered and concentrated under reduced pressure. The residue was taken up in CHCl₃ (200 ml) and washed with water. The organic layer was dried (Na₂SO₄) and concentrated under reduced pressure to give a yellow-brown oil. The crude product was purified by column chromatography (silica gel, CH₂Cl₂-EtOH 98:2). The fractions containing pure product were collected and dried to give an limpid oil (1.09 g; 1.68 mmol; 37%). ¹H NMR (CDCl₃): d 1.13 (m, 4H), 1.27 (t, 12H), 1.73 (m, 2H), 1.86 (m, 2H), 2.30-2.85 (m, 4H+2H), 2.95 (m, 2H), 3.55 (s, 8H), 4.17 (m, 12H). ¹³C NMR (CDCl₃): d 14.25, 18.63, 25.79, 27.24, 47.35, 48.98, 53.78, 55.43, 60.54, 62.64, 119.49, 171.11. (M+H⁺): 652

trans-cyclohexane-1,2-diamine-*N,N'*-propionic-*N,N'*-[*N''*,*N''*'-bis-(2-aminoethyl)]-tetra-acetic acid 4d: this hexaacid has been prepared as described above for compounds 4a & 4b from 1 g of the ester 3d and two volumes of 20 ml HCl (6N). ¹H NMR (D₂O): 1.20-2.00 (m, 10 H), 2.30 (m, 2H), 2.40-3.00 (m, 4H), 3.10-3.95 (m, 14 H), 4.20 (m, 4H) d ¹³C NMR (D₂O) : d 26.89, 30.49, 40.99, 52.00, 55.18, 55.47, 58.97, 165.50, 170.38. (M-H⁺): 547

N,N'-(diethylphosphono-methyl)-*trans*-cyclohexane-1,2-diamine 7: was synthesised in our laboratory according the synthesis procedure of Baily and Burgada with minor modifications. Freshly distilled *trans*-1,2-diaminocyclohexane 1 (3.6 ml, 30 mmol) and diethyl phosphite (7.24 ml, 60 mmol) were dissolved in THF (40ml). The mixture was stirred at reflux and paraformaldehyde (2.8 g, 93 mmol) was added over a 30-min period and the reaction mixture was stirred at reflux for 4 hours. The solvent was evaporated to afford a residue which was taken up in CHCl₃. The organic layer was washed with brine (2*100 ml), dried and evaporated to leave a crude oil. A purification by column chromatography (silica gel, CH₂Cl₂-EtOH 96:4) gave 6 (9.2 g, 21.60 mmol, 72%). 6 was then dissolved in MeOH (40 ml) and 35% HCl (15 ml) was added. The mixture was stirred at 50°C overnight. MeOH was removed, the aqueous layer was adjusted to 50 ml with H₂O and then neutralized by HNaCO₃. bisphosphonate 7 was extracted by CHCl₃. Organic layers were collected, dried and evaporated to give 5.6 g of bisphosphonate 7 (13.52 mmol, 62%, 45% overall). ¹H NMR (CDCl₃): 1.10 (m, 2 H), 1.27 (t, 12H), 1.48 (m, 2H), 1.76 (m, 2 H), 2.13 (m, 2H), 2.98 (m, 2H), 3.10 (t, 2H), 3.35 (t, 2H), 4.12 (m, 8H). ¹³C NMR (CDCl₃): d: 16.36, 16.39, 16.45, 16.38, 24.01, 28.23, 39.67 (JP-C = 156 Hz), 60.74, 60.89, 63.16, 63.23, 63.26, 63.33. (M+H⁺): 415

N,N'-(diethylphosphono-methyl)-*N*-[*N''*,*N''*-bis-(ethylacetate-2-aminoethyl)]-*trans*-cyclohexane-1,2-diamine 8: The mixed ester has been prepared in a mixed solvent system (CH₃CN-H₂O, 1:1) as described above for compound 3d from bisphosphonate 6 (1 g, 2.41 mmol), Na₂HPO₄ (0.5 g; 3.50 mmol) and 1 equivalent of bromide 5 (0.71 g; 2.41 mmol). After stirring for 2 days at 70°C, the reaction mixture was filtered and concentrated under reduced pressure. The residue was taken up in CHCl₃ (200 ml) and washed with water. The organic layer was dried (Na₂SO₄) and concentrated under reduced pressure to give a yellow-brown oil which was used directly in the next step. ¹H NMR: 1.00-1.20 (m, 4 H), 1.25 (t, 6H), 1.31 (t, 12H), 1.72 (m, 2H), 2.04 (m, 2H), 2.23 (m, 1 H), 2.70-2.95 (m, 6H), 3.12 (m, 2H), 3.65 (s, 4H), 4.15 (m, 18H).

N,N'-(diethylphosphono-methyl)-*N'*-(ethylacetate)-*N*-[*N''*,*N''*-bis-(ethylacetate-2-aminoethyl)]-*trans*-cyclohexane-1,2-diamine 9: The tetraester is prepared as described above for compounds 3a & 3b from compound 8,

Na₂CO₃, KI and ethylbromoacetate (1 equivalent). Purification by chromatography (SiO₂) gave a oil.

N,N'-(diethylphosphono-methyl)-*N,N'*-[*N'',N''*-bis-(ethylacetate-2-aminoethyl)]-*trans*-cyclohexane-1,2-diamine 12: The mixed ester is prepared as described above for compound 3d from bisphosphonate 7 (1g, 2.41 mmol), Na₂HPO₄ (1 g, 7.04 mmol) and 2 equivalents of bromide 5 1.80 g, 6.08 mmol). Purification by chromatography (SiO₂ (CH₂Cl₂-MeOH 95 : 5) gave a pale yellow oil (0.59 g, 0.70 mmol, 29%). ¹H NMR: 1.16 (m, 4H), 1.25 (t, 12H), 1.33 (t, 12H), 1.69 (m, 2H), 1.86 (m, 2H + 2H), 2.70-3.50 (m, 8H + 2H + 2H), 3.57 (s, 8H), 4.12 (m, 16H). (M+H⁺): 847



trans-cyclohexane-1,2-diamine-*N,N'*-methylphosphonic-*N*-acetic-,*N'*-[*N'',N''*-bis-(2-aminoethyl)]-tetra-acetic acid 10 & *trans*-cyclohexane-1,2-diamine-*N,N'*-methylphosphonic-*N,N'*-[*N'',N''*-bis-(2-aminoethyl)]-tetra-acetic acid 13 : Those compounds are prepared as described above for compound 4a, 4b and 4d.

Compound 13: ¹H NMR (D₂O): 1.26 (m, 2 H), 1.45 (m, 2H), 1.75-2.00 (m, 4H), 2.80 (m, 2H), 3.00-3.70 (m, 12H), 4.10 (br s, 8H). (M-H⁺): 619

trans-cyclohexane-1,2-diamine-*N,N,N'*-methylphosphonic-*N'*-[*N'',N''*-bis-(2-aminoethyl)]-di-methylphosphonic acid 11 & *trans*-cyclohexane-1,2-diamine-*N,N'*-methylphosphonic-*N,N'*-[*N'',N''*-bis-(2-aminoethyl)]-tetra-methylphosphonic acid 14 : Those compounds are prepared as described above for compound 4c.

153Sm complexation studies

Complexation studies with Samarium 153 were performed on the two following chelating agents 10 (AL 247) and 13 (AL 245)



Radiochemistry purity was measured on ITLC-SG chromatographic profiles. Radioactivity was quantified using a Phosphorimager 445SI apparatus.

Samarium 153 was furnished under $^{153}\text{SmCl}_3$ form in HCl 0,04N with a 5,2 GBq/ml volumic activity and a 40 GBq/mg specific activity.

5 They were tested for their complexation properties by using an excess of 10 to 50 equivalents of chelating agent.

Competition studies were performed against EDTMP according to the following method :

10 - first step : 50 equivalents of one of the chelating agents and a fixed amount of ^{153}Sm were incubated at 37°C during 3 hours in order to form the ^{153}Sm -CA complex (CA = Chelating Agent),

15 - second step : 50 equivalents of EDTMP were added to the previous solution and kept 3h at 37°C in order to measure the decomplexation. Another measure was performed 72h after to ensure a complete decomplexation possibility.

Results :

% of non decomplexed ^{153}Sm -CA		
chelating agent	after 3h	after 72h
10 (AL 247)	100	67
20 13 (AL 245)	100	100

AL 247 and AL 245 present very good complexation properties for ^{153}Sm and in any cases better than EDTMP


25 Furthermore, stability was performed on AL 245 in human serum media at 37°C at different time and showed no loss of ^{153}Sm from AL 245 at either 24, 48, 72 and 96h.

References

- T. Baily and R. Burgada, *Phosphorus, Sulfur and Silicon.*, 1995, 101, 131.
- 5 - M. Bardies, P. Thedrez, J.F. Gestin, B.M. Marcille, D. Guerreau, A. Faivre-Chauvet, M. Mahé, C. Sai-Maurel and J.F. Chatal, *Int. J. Cancer*, 1992, 50, 984.
- R. J. Bergeron, P.S. Burton, K.A. McGovern and S.J. Kline, *Synthesis*, 1981, 732
- 10 - J-F. Gestin, E. Benoist, A. Loussouarn, A.K. Mishra, A. Faivre-Chauvet and J-F. Chatal, *New J. of Chem.*, 1997, 21, 1021.
- W. F. Goeckeler, B. Edwards, W.A. Volkert, R.A. Holmes, J. Simon and D. Wilson, *J. Nucl. Med.*, 1987, 28, 495.
- F. Krüger and L. Bauer, *Chem. Ztg.*, 1972, 36, 691.
- 15 - A. Loussouarn, M. Duflos, E. Benoist, J-F. Chatal, G. Le Baut and J-F. Gestin, *J. Chem. Soc. Perkin Trans.*, 1998, 1, 237.
- C.F. Meares, M.J. Mc Call, D.T. Reardan, D.A. Goodwin, C.I. Diamanti and M. McTigue, *Anal. Chem.*, 1984, 142, 68.
- R.C. Mease, S.C. Srivastava, G.E. Meinken, J-F. Gestin and Z. Steplewski, *J. Nucl. Med.*, 1990, 31, 896.
- 20 - P. L. Ornstein, J. M. Schaus, J. W. Chambers, D. L. Huser, J. D. Leander, D. T. Wong, J. W. Paschal, N. D. Jones and J. B. Deeter, *J. Med. Chem.* 1989, 32, 827.
- D. Parker, *Chem. Soc. Review*, 1990, 19, 271.
- 25 - P.A. Schubiger, R. Alberto and A. Smith, *Bioconjugate Chem.*, 1996, 7, 165.
- R. Stein, D. M. Goldenderg, S. R. Thorpe, A. Basu and M. J. Mattes, *Cancer Research*, 1995, 55, 3132.
- M. Studer and C.F. Meares, *Bioconjugate Chem.*, 1992, 3, 420.
- 30 - M. A. Williams and H. Rapoport, *J. Org. Chem.*, 1994, 59, 3616.

CLAIMS

1. Compounds of the following formula (I) :




in which :

- n_1 , n_2 , n_3 and n_4 , independently from each other, represent an integer from 1 to 5, preferably from 1 to 3,

- R_1 , R_2 , R_3 and R_4 , independently from each other, represent :


. $-\text{COOH}$,

. $-\text{PO}(\text{OH})_2$,



wherein n_5 represents an integer from 1 to 5, preferably from 1 to 3, R_5 represents $-\text{COOH}$ or $-\text{PO}(\text{OH})_2$, and Y represents H or a group $-(\text{CH}_2)_{n_6} - \text{R}_6$ in which n_6 represents an integer from 1 to 5, preferably from 1 to 3, and R_6 represents $-\text{COOH}$ or $-\text{PO}(\text{OH})_2$,

provided that at least one of R_1 , R_2 , R_3 or R_4 represents a group



such as defined above,



- R represents :

. H , or $-\text{NHCOCH}_3$, or

. a group carrying a function liable to bind, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, which are able to bind specifically with epitopes located at the surface of the cells of the organism, or to chemical or biological compounds located at the surface of a solid carrier, or

: a group carrying a function linked, if necessary via a binding site, to molecules as defined above,


the two following compounds, CDTPA and CTTHA, being excluded :



2. Compounds according to claim 1, characterized in that :

- when R₁, R₂, R₃ or R₄ represents -COOH or -PO(OH)₂, then n₁, n₂, n₃ or n₄ represents 1 respectively,


- when R₁, R₂, R₃ or R₄ represents a group



then n₁, n₂, n₃ or n₄ represents 2 or 3 respectively, and preferably 2,

- n₅, and optionally n₆, represents 1.

3. Compounds according to claims 1 or 2, characterized in that at least one, and more preferably two of R₁, R₂, R₃ and R₄, represent a group




wherein n₅, n₆, R₅ and R₆ are defined in claims 1 or 2.

4. Compounds according to anyone of claims 1 to 3, characterized in that R represents a group carrying a function liable to bind, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, as defined in

claim 1, and in particular R represents a group chosen among all the coupling functions for vector or solid support binding.

5 5. Compounds according to anyone of claims 1 to 3, characterized in that R represents a group carrying a function linked, if necessary via a binding site, to molecules, such as antibodies, haptens or peptides, as defined in claim 1.

10 6. Compounds according to anyone of claims 1 to 5 of the following formula (III) :




20 in which R₁, R₂, R₃, R₅ and R₆ independently from each other represent -COOH or -PO(OH)₂, and R is a group as defined in claims 2 to 5.

25 7. Compounds according to claim 6, of formula (III) wherein :

- . R₁ = R₅ = R₆ = COOH and R₂, = R₃ = PO(HO)₂, or
- . R₁ = R₂ = R₃ = R₅ = R₆ = COOH, or
- . R₁ = R₂ = R₃ = R₅ = R₆ = PO(OH)₂.

30 8. Compounds according to anyone of claims 1 to 5, of the following formula (IV) :



35 wherein R₂, R₅ and R₆, independently from each other, represent -COOH or -PO(OH)₂, and R is a group as defined in claims 2 to 5.

9. Compounds according to claim 8 of formula (IV) wherein :

- . $R_2 = R_3 = PO(OH)_2$, and $R_5 = R_6 = COOH$, or
- . $R_2 = R_3 = R_5 = R_6 = COOH$, or
- . $R_2 = R_3 = R_5 = R_6 = PO(OH)_2$.

5

10. Complexes between a compound according to anyone of claims 1 to 9, and a radioactive element.

11. Complexes according to claim 10, characterized in that said radioelements are α or β emitter radiometals.

10

12. Complexes according to claim 11, characterized in that the compound is chosen among those defined in anyone of claims 5 to 9, and more particularly among those compounds wherein the group R comprises :

15

- an antibody (polyclonal or monoclonal) liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism,

- or an hapten, i.e. a non-immunogenic molecule of low MW capable of inducing the production of antibodies against itself, said hapten being liable to recognize and to bind to one or several molecules already bound, in a first step of the treatment, to epitopes on the surface of specific cells in the organism,

20

- or a peptide resulting from the association of different amino acids and liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism.

25

13. Use of a complex according to claims 11 or 12, for the manufacture of a medicament for radioimmunotherapy, such as for the treatment of cancers, and more particularly for the treatment against metastase proliferation.

14. Pharmaceutical compositions characterized in that they comprise an effective amount of a complex according to claims 11 or 12, in association with a suitable pharmaceutical carrier.

30

15. Complexes according to claim 10, characterized in that the radioelements are γ emitter radiometals.

35

16. Complexes according to claim 15, characterized in that the compound is chosen among those defined in anyone of claims 5 to 9, and more particularly among those compounds wherein the group R comprises :

- an antibody (polyclonal or monoclonal) liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism,

- or an hapten, i.e. a non-immunogenic molecule of low MW capable of inducing the production of antibodies against itself, said hapten being liable to recognize and to bind to one or several molecules already bound (in a first step of the method of diagnosis) to epitopes on the surface of specific cells in the organism,

- or a peptide resulting from the association of different amino acids and liable to recognize and to bind to specific epitopes on the surface of specific cells of the organism.

17. Use of a complex according to claims 15 or 16, for carrying out diagnosis methods such as radioimmunoscintigraphy.

18. Use of a compound of formula (I) as defined in claim 1 to 9, included compounds CDTPA and CTTHA, for :

- the manufacture of a medicament useful as antalgic, or for the treatment of pathologies where ionic imbalances occur, or against the formation of stones in the organism,

- carrying out a process for the detoxication of polluted medium, such as liquid phases polluted by bivalent or trivalent metals radioactives or not,

- carrying out a process for the radionuclides purification, said compound being bound to a solid phase,

- carrying out a bone scintigraphy, in particular in the frame of the diagnosis of osteoarticular pathology, particularly in bone cancer extension balance.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/08031

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07F9/38 C07C229/16 A61K51/10 A61K51/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07F C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 292 938 A (RONNIE C. MEASE) 8 March 1994 (1994-03-08) the whole document — —/—	1-18

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

21 December 1999

Date of mailing of the international search report

11/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Beslier, L

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/08031

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>LOUSSOUARN A ET AL: "Synthesis of new bifunctional chelating agents: (1R*,2R*,4S*)-4-isothiocyanatocyclohexane-1,2-diamine-N,N,N',N'-tetra kismethanephosphonic acid (4-ICMP) and (1R*,2R*,4S*)-4-isothiocyanato cyclohexane-1,2-diamine-N,N,N',N'-tetrakis ethanephosphonic acid (4-ICEP)" J. CHEM. SOC., PERKIN TRANS. 1 (JCPRB4,0300922X);1998; (2); PP.237-242, - 21 January 1998 (1998-01-21) XP002097516 Institut de Biologie;INSERM U.463 (ex-U.211); Nantes; 44035; Fr. (FR) cited in the application the whole document</p>	1-18
Y	<p>BRECHBIEL M W ET AL: "Preparation of the Novel Chelating Agent N-(2-Aminoethyl)-trans-1,2 -diaminocyclohexane-N,N',N''-pentaacetic Acid (H5CyDTPA), a Preorganized Analog of Diethylenetriaminepentaacetic Acid (H5DTPA), and the Structures of BiIII(CyDTPA)2- and BiIII(H2DTPA) Complexes" INORG. CHEM. (INOCAL,00201669); VOL.35 (21); PP.6343-6348, - 9 October 1996 (1996-10-09) XP002097517 National Institutes of Health;Chemistry Section; Bethesda; 20892; MD; USA (US) the whole document</p>	1-18
Y	<p>US 5 089 663 A (RONNIE C. MEASE) 18 February 1992 (1992-02-18) the whole document</p>	1-18
P,X	<p>LOUSSOUARN A ET AL: "Simple and general procedure for the synthesis of semi-rigid chelating agents for radiometal complexation studies and its application to semi-rigid functionalised ligands (BCA) synthesis" MATER. SCI. FORUM (MSFOEP,02555476);1999; VOL.315-317 (RARE EARTHS '98); PP.262-267, XP000865647 Institut Biologie;Nantes; F-44035; Fr. (FR) the whole document</p>	1-18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/08031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5292938 A	08-03-1994	US 5635157 A	03-06-1997
US 5089663 A	18-02-1992	US 5021571 A	04-06-1991
		US 5334729 A	02-08-1994